arXiv:1512.00395v1 [nlin.SI] 1 Dec 2015

A non autonomous generalization of the $Q_{\rm V}$ equation

G. Gubbiotti*, C. Scimiterna†, D. Levi‡
Dipartimento di Matematica e Fisica, Universita' degli Studi Roma Tre,
e Sezione INFN di Roma Tre,
Via della Vasca Navale 84, 00146 Roma (Italy)

December 2, 2015

Abstract

In this paper we introduce a non autonomous generalization of the $Q_{\rm V}$ equation introduced by Viallet. All the equations of Boll's classification appear in it for special choices of the parameters. Using the algebraic entropy test we infer that the equation should be integrable and with the aid of a formula introduced by Xenitidis we find its three point generalized symmetries.

1 Introduction

In a previous paper [10] we calculated the three point generalized symmetries of equations belonging to Boll classification [4–6] of quad graph equations compatible around the cube, i.e. the trapezoidal H^4 equations and the H^6 equations. The symmetres of the rhombic H^4 equations has been considered in [27]. In [10] we also noticed that all the fluxes of such symmetries are related to some particular cases of the non autonomous YdKN equation [17][1]:

$$\frac{\mathrm{d}u_n}{\mathrm{d}t} = \frac{A_n(u_n)u_{n+1}u_{n-1} + B_n(u_n)(u_{n+1}+u_{n-1}) + C_n(u_n)}{u_{n+1}-u_{n-1}}. \tag{1}$$

Here the n-dependent coefficients are given by:

$$A_n(u_n) = au_n^2 + 2b_nu_n + c_n, \tag{2a}$$
$$B_n(u_n) = b_{n+1}u_n^2 + du_n + e_{n+1}, \tag{2b}$$
$$C_n(u_n) = c_{n+1}u_n^2 + 2e_nu_n + f, \tag{2c}$$

where b_n, c_n and e_n are 2-periodic functions, i.e.

$$b_n = b_0 + b_1(-1)^n,\ c_n = c_0 + c_1(-1)^n,\ e_n = e_0 + e_1(-1)^n. \tag{3}$$

*e-mail: gubbiotti@mat.uniroma3.it
†e-mail: scimiterna@fis.uniroma3.it
‡e-mail: decio.levi@roma3.infn.it

[1]For the rhombic H^4 equations this was already known from [27].

[10] naturally extended the results contained in [14] where the three points generalized symmetries of the ABS class of lattice equations [1], which are at the base of Boll's classification, are shown to be sub–cases of the autonomous YdKN equation [28]. We recall that the autonomous YdKN equation is obtained from (1) just taking b_k, c_k and e_k as pure constants, i.e.:

$$\frac{\mathrm{d}u_n}{\mathrm{d}t} = \frac{A(u_n)u_{n+1}u_{n-1} + B(u_n)(u_{n+1} + u_{n-1}) + C(u_n)}{u_{n+1} - u_{n-1}}. \tag{4}$$

where:

$$A(u_n) = au_n^2 + 2bu_n + c, \tag{5a}$$
$$B(u_n) = bu_n^2 + du_n + e, \tag{5b}$$
$$C(u_n) = cu_n^2 + 2eu_n + f, \tag{5c}$$

In particular in [10] we noted that all equations in the Boll classification are subcases of the general non autonomous YdKN (1) corresponding to $a = b_k = 0$.

In [26] it was shown that the Q_{V} equation, introduced in [24]:

$$\begin{aligned} Q_{\mathrm{V}} =& a_1 u_{n,m}u_{n+1,m}u_{n,m+1}u_{n+1,m+1} \\ &+a_{2,0}\left(u_{n,m}u_{n,m+1}u_{n+1,m+1} + u_{n+1,m}u_{n,m+1}u_{n+1,m+1}\right. \\ &\quad \left.+u_{n,m}u_{n+1,m}u_{n+1,m+1} + u_{n,m}u_{n+1,m}u_{n,m+1}\right) \\ &+a_{3,0}\left(u_{n,m}u_{n+1,m} + u_{n,m+1}u_{n+1,m+1}\right) \\ &+a_{4,0}\left(u_{n,m}u_{n+1,m+1} + u_{n+1,m}u_{n,m+1}\right) \\ &+a_{5,0}\left(u_{n+1,m}u_{n+1,m+1} + u_{n,m}u_{n,m+1}\right) \\ &+a_{6,0}\left(u_{n,m} + u_{n+1,m} + u_{n,m+1} + u_{n+1,m+1}\right) \\ &+a_7 = 0 \end{aligned} \tag{6}$$

admits a symmetry in the direction n

$$\frac{du_{n,m}}{dt} = \frac{h_n}{u_{n+1,m} - u_{n-1,m}} - \frac{1}{2}\partial_{u_{n+1,m}} h_n. \tag{7}$$

where:

$$\begin{aligned} h_n(u_{n,m}, u_{n+1,m}) =& Q_V \partial_{u_{n,m+1}}\partial_{u_{n+1,m+1}} Q_V \\ &- \left(\partial_{u_{n,m+1}} Q_V\right)\left(\partial_{u_{n+1,m+1}} Q_V\right) \end{aligned} \tag{8}$$

and a symmetry in the direction m

$$\frac{du_{n,m}}{dt} = \frac{h_m}{u_{n,m+1} - u_{n,m-1}} - \frac{1}{2}\partial_{u_{n,m+1}} h_m. \tag{9}$$

where:

$$\begin{aligned} h_m(u_{n,m}, u_{n,m+1}) =& Q_V \partial_{u_{n+1,m}}\partial_{u_{n+1,m+1}} Q_V \\ &- \left(\partial_{u_{n+1,m}} Q_V\right)\left(\partial_{u_{n+1,m+1}} Q_V\right) \end{aligned} \tag{10}$$

of the same form as the YdKN (4).

The connection formulae between the coefficient of Q_V and the n direction YdKN (4) are:

$$a = a_{3,0}a_1 - a_{2,0}^2, \qquad b = \frac{1}{2}[a_{2,0}(a_{3,0} - a_{5,0} - a_{4,0}) + a_{6,0}a_1],$$

$$c = a_{2,0}a_{6,0} - a_{4,0}a_{5,0}, \qquad d = \frac{1}{2}[a_{3,0}^2 - a_{4,0}^2 - a_{5,0}^2 + a_1 a_7], \tag{11}$$
$$e = \frac{1}{2}[a_{6,0}(a_{3,0} - a_{4,0} - a_{5,0}) + a_{2,0}a_7], \qquad f = a_{3,0}a_7 - a_{6,0}^2.$$

and show that (a, b) may be different from $(0, 0)$. The connection formulae between the coefficient of Q_V and the m direction YdKN (4) are:

$$\begin{aligned} a &= a_{5,0}a_1 - a_{2,0}^2, \qquad b = \frac{1}{2}[a_{2,0}(a_{5,0} - a_{3,0} - a_{4,0}) + a_{6,0}a_1], \\ c &= a_{2,0}a_{6,0} - a_{4,0}a_{3,0}, \qquad d = \frac{1}{2}[a_{5,0}^2 - a_{4,0}^2 - a_{3,0}^2 + a_1 a_7], \\ e &= \frac{1}{2}[a_{6,0}(a_{5,0} - a_{4,0} - a_{3,0}) + a_{2,0}a_7], \qquad f = a_{5,0}a_7 - a_{6,0}^2. \end{aligned} \tag{12}$$

So the three point generalized symmetries of the $Q_{\rm V}$ belong to the class of the general YdKN equation (4).

From the results obtained in [10] we were led to conjecture the existence of a non autonomous generalization of the $Q_{\rm V}$ equation and suggested some possible ways to obtain such a generalization. In this paper we follow them to generalize the $Q_{\rm V}$ equation. It is known [24] that $Q_{\rm V}$ is the most general multi-linear equation on a quad graph possessing Klein discrete symmetries, i.e. such that:

$$\begin{aligned} Q\left(u_{n+1,m}, u_{n,m}, u_{n+1,m+1}, u_{n,m+1}\right) &= \tau Q\left(u_{n,m}, u_{n+1,m}, u_{n,m+1}, u_{n+1,m+1}\right), \\ Q\left(u_{n,m+1}, u_{n+1,m+1}, u_{n,m}, u_{n,m+1}\right) &= \tau' Q\left(u_{n,m}, u_{n+1,m}, u_{n,m+1}, u_{n+1,m+1}\right), \end{aligned} \tag{13}$$

where $\tau, \tau' = \pm 1$. A way to extend the Klein discrete symmetry is to consider a multi-linear function Q with two-periodic coefficients in n and m such that the following equations holds:

$$\begin{aligned} &Q\left(u_{n+1,m}, u_{n,m}, u_{n+1,m+1}, u_{n,m+1}; (-1)^n, (-1)^m\right) = \\ &\quad \tau Q\left(u_{n,m}, u_{n+1,m}, u_{n,m+1}, u_{n+1,m+1}; -(-1)^n, (-1)^m\right), \\ &Q\left(u_{n,m+1}, u_{n+1,m+1}, u_{n,m}, u_{n+1,m}; (-1)^n, (-1)^m\right) = \\ &\quad \tau' Q\left(u_{n,m}, u_{n+1,m}, u_{n,m+1}, u_{n+1,m+1}; (-1)^n, -(-1)^m\right). \end{aligned} \tag{14}$$

If a non autonomous system Q satisfies the discrete symmetries (14) we will say that Q admits a *non autonomous discrete Klein symmetry*. The name follows from the fact that if Q is autonomous then the discrete symmetry (14) reduces to the Klein one (13). Furthermore all the equation belonging to the Boll's classification satisfy this symmetry conditions (14) when $\tau = \tau' = 1$.

In Section 2 we will show that up to multiplication by a function and re-definition of the parameters there is only one quad graph equation possessing the non autonomous Klein symmetry. We will prove that this equation, which we will call the *non autonomous Q_V equation* has as sub–cases all the equations of Boll classification and it satisfies the Algebraic Entropy integrability test. Furthermore we will show that (7, 8) will provide an n directional and (9, 10) an m directional symmetry for the non autonomous $Q_{\rm V}$ equation and that such symmetries belong to the class of the non autonomous YdKN equation (1, 2). We will provide then the appropriate connection formulæ. In Section 3 we will present some concluding remarks.

2 The non autonomous $Q_{\rm V}$ equation

Let us consider the most general multi-linear equation in the lattice variables with two-periodic coefficients:

$$\begin{aligned}
&p_1 u_{n,m}u_{n+1,m}u_{n,m+1}u_{n+1,m+1}\\
&+p_2 u_{n,m}u_{n,m+1}u_{n+1,m+1}+p_3 u_{n+1,m}u_{n,m+1}u_{n+1,m+1}\\
&+p_4 u_{n,m}u_{n+1,m}u_{n+1,m+1}+p_5 u_{n,m}u_{n+1,m}u_{n,m+1}\\
&+p_6 u_{n,m}u_{n+1,m}+p_7 u_{n,m+1}u_{n+1,m+1}+p_8 u_{n,m}u_{n+1,m+1}\\
&+p_9 u_{n+1,m}u_{n,m+1}+p_{10} u_{n+1,m}u_{n+1,m+1}+p_{11} u_{n,m}u_{n,m+1}\\
&+p_{12} u_{n,m}+p_{13} u_{n+1,m}+p_{14} u_{n,m+1}+p_{15} u_{n+1,m+1}+p_{16}=0
\end{aligned} \tag{15}$$

i.e. the p_i coefficients have the following expression:

$$p_i = p_{i,0}+p_{i,1}(-1)^n+p_{i,2}(-1)^m+p_{i,3}(-1)^{n+m}, \quad i=1,\dots,16. \tag{16}$$

If we impose the non autonomous Klein symmetry condition (14) with $\tau=\tau'=1$ the 64 coefficients of (15) turn out to be related among themselves and we can choose among them 16 independent coefficients. In term of the 16 independent coefficients (15) reads:

$$\begin{aligned}
&a_1 u_{n,m}u_{n+1,m}u_{n,m+1}u_{n+1,m+1}\\
&+\left[a_{2,0}-(-1)^n a_{2,1}-(-1)^m a_{2,2}+(-1)^{n+m}a_{2,3}\right]u_{n,m}u_{n,m+1}u_{n+1,m+1}\\
&+\left[a_{2,0}+(-1)^n a_{2,1}-(-1)^m a_{2,2}-(-1)^{n+m}a_{2,3}\right]u_{n+1,m}u_{n,m+1}u_{n+1,m+1}\\
&+\left[a_{2,0}+(-1)^n a_{2,1}+(-1)^m a_{2,2}+(-1)^{n+m}a_{2,3}\right]u_{n,m}u_{n+1,m}u_{n+1,m+1}\\
&+\left[a_{2,0}-(-1)^n a_{2,1}+(-1)^m a_{2,2}-(-1)^{n+m}a_{2,3}\right]u_{n,m}u_{n+1,m}u_{n,m+1}\\
&+\left[a_{3,0}-(-1)^m a_{3,2}\right]u_{n,m}u_{n+1,m}\\
&+\left[a_{3,0}+(-1)^m a_{3,2}\right]u_{n,m+1}u_{n+1,m+1}\\
&+\left[a_{4,0}-(-1)^{n+m}a_{4,3}\right]u_{n,m}u_{n+1,m+1}\\
&+\left[a_{4,0}+(-1)^{n+m}a_{4,3}\right]u_{n+1,m}u_{n,m+1}\\
&+\left[a_{5,0}-(-1)^n a_{5,1}\right]u_{n+1,m}u_{n+1,m+1}\\
&+\left[a_{5,0}+(-1)^n a_{5,1}\right]u_{n,m}u_{n,m+1}\\
&+\left[a_{6,0}+(-1)^n a_{6,1}-(-1)^m a_{6,2}-(-1)^{n+m}a_{6,3}\right]u_{n,m}\\
&+\left[a_{6,0}-(-1)^n a_{6,1}-(-1)^m a_{6,2}+(-1)^{n+m}a_{6,3}\right]u_{n+1,m}\\
&+\left[a_{6,0}+(-1)^n a_{6,1}+(-1)^m a_{6,2}+(-1)^{n+m}a_{6,3}\right]u_{n,m+1}\\
&+\left[a_{6,0}-(-1)^n a_{6,1}+(-1)^m a_{6,2}-(-1)^{n+m}a_{6,3}\right]u_{n+1,m+1}\\
&+a_7=0
\end{aligned} \tag{17}$$

Upon the substitution $a_{2,1}=a_{2,2}=a_{2,3}=a_{3,2}=a_{4,3}=a_{5,1}=a_{6,1}=a_{6,2}=a_{6,3}=0$ (17) reduces to the $Q_{\rm V}$ equation (6). Therefore we will call (17) *the non autonomous Q_V equation.*

If we impose the non autonomous Klein symmetry condition (14) with the choice $\tau = 1$ and $\tau' = -1$ we will get an expression which can be reduced to (17) by multiplying by $(-1)^n$ and redefining the coefficients. In an analogous manner the two remaining cases $\tau = -1$, $\tau' = 1$ and $\tau = \tau' = -1$ can be identified with the case $\tau = \tau' = 1$ multiplying by $(-1)^n$ and $(-1)^{n+m}$ respectively and redefining the coefficients. Therefore the only equation belonging to the class of the lattice equation possessing the non autonomous Klein symmetries is just the non autonomous $Q_{\rm V}$ equation (17).

We note that the non autonomous $Q_{\rm V}$ equation contains as particular cases the rhombic H^4 equations, the trapezoidal H^4 equations and the H^6 equations. The explicit identification of the coefficients of such equations is given in Table 1. The reader can refer to Appendix A for the explicit expressions of these equations or to [9] for a complete derivation following the prescription of [6].

In order to establish if (17) is integrable we use the Algebraic Entropy integrability test [19, 22, 23], using the program `ae2d.py` [8, 18]. Applying it to the non autonomous $Q_{\rm V}$ equation we find the following degree of growth in all directions:

$$1, 3, 7, 13, 21, 31, 43, 57, 73, 91, 111, 133 \ldots, \tag{18}$$

which is the same as for the autonomous $Q_{\rm V}$ equation [24]. The generating function for the sequence (18) is:

$$g(z) = \frac{1 + z^2}{(1 - z)^3}, \tag{19}$$

which implies that we have the following quadratic fit for the growth:

$$d_k = k(k + 1) + 1, \tag{20}$$

and thus the Algebraic Entropy is zero. This is a strong indication of the integrability of the non autonomous $Q_{\rm V}$ equation (17).

Using (7, 8) or (9, 10) with $Q_{\rm V}$ substituted by its non autonomous version we get a version of the non autonomous YdKN (1) however the proof that this is effectively a symmetry of the non autonomous $Q_{\rm V}$ encounters serious computational difficulties.

We can prove by a direct computation its validity for the following sub-cases:

- When Qv equation is non autonomous with respect to one direction only, either n or m. All the trapezoidal ${}_tH^4$ equations belong to these two sub-classes;
- For all the H^6 equations, which are non autonomous in both directions.

Its validity for the autonomous Qv and for all the rhombic ${}_rH^4$ equations was already showed respectively in [26] and [27]. However we cannot prove its validity for the general case (17).

Here in the following we compute the connection formulae for the general non autonomous case (17). For the n directional symmetry we have:

$$\begin{aligned} a &= a_1 a_{3,0} - a_{2,0}^2 + \\ &\quad + a_{2,1}^2 - a_{2,2}^2 + a_{2,3}^2 - (-1)^m (2a_{2,0}a_{2,2} - 2a_{2,1}a_{2,3} + a_1 a_{3,2}), \\ b_0 &= \frac{1}{2}\{a_{2,0}(a_{3,0} - a_{5,0} - a_{4,0}) + a_1 a_{6,0} + a_{2,2}a_{3,2} - a_{2,3}a_{4,3} - a_{2,1}a_{5,1} - \end{aligned} \tag{21}$$

$$
\begin{aligned}
& -(-1)^m[a_{2,2}(a_{5,0}+a_{3,0}+a_{4,0})+a_{2,3}a_{5,1}+a_1a_{6,2}+a_{2,0}a_{3,2}+ \\
& +a_{2,1}a_{4,3}]\},
\end{aligned}
$$

$$
\begin{aligned}
b_1 &= \frac{1}{2}\{a_{2,1}(a_{3,0}-a_{4,0}+a_{5,0})+a_{2,3}a_{3,2}-a_{2,2}a_{4,3}+a_{2,0}a_{5,1}-a_1a_{6,1}+ \\
&\quad +(-1)^m[a_1a_{6,3}-a_{2,3}(a_{3,0}+a_{4,0}-a_{5,0})-a_{2,1}a_{3,2}-a_{2,0}a_{4,3}+ \\
&\quad +a_{2,2}a_{5,1}]\},
\end{aligned}
$$

$$
\begin{aligned}
c_0 &= a_{2,0}a_{6,0}-a_{4,0}a_{5,0}-a_{2,1}a_{6,1}-a_{2,3}a_{6,3}+a_{2,2}a_{6,2}+ \\
&\quad -(-1)^m[a_{2,2}a_{6,0}-a_{4,3}a_{5,1}-a_{2,3}a_{6,1}+a_{2,0}a_{6,2}-a_{2,1}a_{6,3}],
\end{aligned}
$$

$$
\begin{aligned}
c_1 &= a_{4,0}a_{5,1}+a_{2,1}a_{6,0}-a_{2,0}a_{6,1}+a_{2,3}a_{6,2}-a_{2,2}a_{6,3}+ \\
&\quad +(-1)^m[a_{2,2}a_{6,1}-a_{4,3}a_{5,0}-a_{2,3}a_{6,0}-a_{2,1}a_{6,2}+a_{2,0}a_{6,3}],
\end{aligned}
$$

$$
\begin{aligned}
d &= \frac{1}{2}[a_{3,0}^2-a_{4,0}^2-a_{5,0}^2+a_1a_7-a_{3,2}^2+a_{4,3}^2+a_{5,1}^2- \\
&\quad -4(-1)^m(a_{2,2}a_{6,0}+a_{2,3}a_{6,1}+a_{2,0}a_{6,2}+a_{2,1}a_{6,3}],
\end{aligned}
$$

$$
\begin{aligned}
e_0 &= \frac{1}{2}\{a_{6,0}[a_{3,0}-a_{4,0}-a_{5,0}]+a_{2,0}a_7+a_{5,1}a_{6,1}-a_{3,2}a_{6,2}+a_{4,3}a_{6,3}+ \\
&\quad +(-1)^m[a_{3,2}a_{6,0}+a_{4,3}a_{6,1}+a_{5,1}a_{6,3}-a_{6,2}(a_{3,0}+a_{4,0}+a_{5,0})-a_{2,2}a_7]\},
\end{aligned}
$$

$$
\begin{aligned}
e_1 &= \frac{1}{2}\{a_{6,1}[a_{3,0}-a_{4,0}+a_{5,0}]-a_{5,1}a_{6,0}+a_{4,3}a_{6,2}-a_{3,2}a_{6,3}-a_{2,1}a_7+ \\
&\quad +(-1)^m[a_{4,3}a_{6,0}+a_{3,2}a_{6,1}-a_{5,1}a_{6,2}+a_{6,3}(a_{5,0}-a_{3,0}-a_{4,0})+a_{2,3}a_7\},
\end{aligned}
$$

$$
f = a_{3,0}a_7-a_{6,0}^2-a_{6,2}^2+a_{6,3}^2+a_{6,1}^2-(-1)^m(2a_{6,0}a_{6,2}-2a_{6,1}a_{6,3}-a_{3,2}a_7).
$$

The non autonomous $Q_{\rm V}$ is not symmetric in the exchange of n and m so its symmetries in the m direction are different for their dependence on the coefficients and so are the connection formulae, however, for the sake of the reader as they are not essentially different from those presente above we do not write them here but present them in Appendix B, cfr. (29).

Eq.	$a_{3,0}$	$a_{3,2}$	$a_{4,0}$	$a_{4,3}$	$a_{5,0}$	$a_{5,1}$	$a_{6,0}$	$a_{6,1}$	$a_{6,2}$	$a_{6,3}$	a_7
${}_rH_1^\varepsilon$	1	0	$\frac{1}{2}\varepsilon(\alpha-\beta)$	$\frac{1}{2}\varepsilon(\alpha-\beta)$	-1	0	0	0	0	0	$\beta-\alpha$
${}_rH_2^\varepsilon$	1	0	$2\varepsilon\,(\beta-\alpha)$	$2\varepsilon\,(\beta-\alpha)$	-1	0	$-(\alpha-\beta)\,(\varepsilon\alpha+1+\varepsilon\beta)$	0	0	$\varepsilon\left(\beta^2-\alpha^2\right)$	$-(\alpha-\beta)\left(2\varepsilon\alpha^2+\alpha+2\varepsilon\beta^2+\beta\right)$
${}_rH_3^\varepsilon$	α	0	$\frac{1}{2}\frac{\varepsilon\left(\beta^2-\alpha^2\right)}{\alpha\beta}$	$\frac{1}{2}\frac{\varepsilon\left(\beta^2-\alpha^2\right)}{\alpha\beta}$	$-\beta$	0	0	0	0	0	$\delta\left(\alpha^2-\beta^2\right)$
${}_tH_1^\varepsilon$	$-\frac{1}{2}\alpha_2\varepsilon^2$	$-\frac{1}{2}\alpha_2\varepsilon^2$	-1	0	1	0	0	0	0	0	$-\alpha_2$
${}_tH_2^\varepsilon$	$\varepsilon\alpha_2$	$\varepsilon\alpha_2$	-1	0	1	0	$\frac{1}{2}\alpha_2\left[2+\varepsilon(2\alpha_2+\alpha_3)\right]$	0	$\varepsilon\alpha_2\alpha_3+\frac{1}{2}\varepsilon{\alpha_2}^2$	0	$\alpha_2\left[\alpha_2+2\alpha_3+\varepsilon\left(\alpha_2+\alpha_3\right)^2\right]$
${}_tH_3^\varepsilon$	$\frac{1}{2}\frac{\varepsilon^2\left(1-{\alpha_2}^2\right)}{\alpha_3\alpha_2}$	$\frac{1}{2}\frac{\varepsilon^2\left(1-{\alpha_2}^2\right)}{\alpha_3\alpha_2}$	α_2	0	-1	0	0	0	0	0	$\delta^2\alpha_3\left(1-\alpha_2^2\right)$
${}_1D_2$	$\frac{1}{2}\delta_1$	$\frac{1}{2}\delta_1$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{1}{2}-\frac{1}{4}(\delta_1-\delta_2)$	$\frac{1}{4}(\delta_2-\delta_1)$	$-\frac{1}{4}(\delta_1+\delta_2)$	$\frac{1}{2}-\frac{1}{4}(\delta_1+\delta_2)$	0
${}_2D_2$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}\delta_1$	$\frac{1}{2}\delta_1$	0	0	$\frac{1}{2}-\frac{1}{4}(\delta_1-\delta_2+\delta_1\lambda)$	$\frac{1}{4}(\delta_1\lambda-\delta_1+\delta_2)$	$\frac{1}{2}-\frac{1}{4}(\delta_1-\delta_1\lambda+\delta_2)$	$-\frac{1}{4}(\delta_1+\delta_1\lambda+\delta_2)$	$-\delta_1\delta_2\lambda$
${}_3D_2$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{1}{2}\delta_1$	$-\frac{1}{2}\delta_1$	$\frac{1}{2}-\frac{1}{4}(\delta_1-\delta_2+\delta_1\lambda)$	$\frac{1}{4}(\delta_1+\delta_1\lambda+\delta_2)$	$\frac{1}{2}-\frac{1}{4}(\delta_1-\delta_1\lambda+\delta_2)$	$\frac{1}{4}(\delta_1-\delta_1\lambda-\delta_2)$	$-\delta_1\delta_2\lambda$
D_3	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	0
${}_1D_4$	$\frac{1}{2}\delta_2$	$\frac{1}{2}\delta_2$	1	0	$\frac{1}{2}\delta_1$	$-\frac{1}{2}\delta_1$	0	0	0	0	δ_3
${}_2D_4$	1	0	$\frac{1}{2}\delta_2$	$\frac{1}{2}\delta_2$	$\frac{1}{2}\delta_1$	$-\frac{1}{2}\delta_1$	0	0	0	0	δ_3

Table 1: Identification of the coefficients of the non autonomous $Q_{\rm V}$ equation with those of the Boll's equations (23,24, 25). Since $a_1 = a_{2,i} = 0$ for every equation these coefficients absent in the Table.

3 Conclusions

In this article we propose a non autonomous extension of the $Q_{\rm V}$ equation (17) which satisfies an extended Klein symmetry (14) in such a way that in the autonomous sub-case reduces to the $Q_{\rm V}$ equation (6). The so obtained non autonomous $Q_{\rm V}$ equation includes all equations of the Boll classification as its sub-cases and results integrable by the Algebraic Entropy test. Using the construction proposed by Xenitidis [26] one builds up a symmetry for the QV non autonomous with respect to one direction only. This symmetry turns out to belong to the class of the non autonomous YdKN equation proposed by Levi and Yamilov [16], an equation satisfying all five integrability conditions obtained by the formal generalized symmetry method. One is confident that the results obtained using the formulae (7, 9) by Xenitidis [26] are correct also in general setting, as for all the sub-cases of the non autonomous QV belonging to the Boll classification the three point generalized symmetries, calculated using both the definition or through the formula by Xenitidis, coincide and belong to the class of the non autonomous YdKN. However we have not been able to have a complete direct proof in the general non autonomous case due to computational complexities and work on it is in progress. In all generality we have two possible connection formulae (21, 29) between the 16 coefficients of the general non autonomous QV and the 9 coefficients of the non autonomous YdKN, one corresponding to the Xenitidis formula (7) along the n direction and another one corresponding to (9) along m.

Acknowledgment

CS and DL have been partly supported by the Italian Ministry of Education and Research, 2010 PRIN *Continuous and discrete nonlinear integrable evolutions: from water waves to symplectic maps.*
GG and DL are supported by INFN IS-CSN4 *Mathematical Methods of Nonlinear Physics.*

A Explicit form the H^4 and H^6 equations

Throughout this appendix we will use the notation:

$$F_k^{\pm} = \frac{1 \pm (-1)^k}{2}, \quad k \in \mathbb{Z}. \tag{22}$$

Rhombic H^4 equations:

$$\begin{aligned} {}_rH_1^{\varepsilon}: \quad & (u_{n,m} - u_{n+1,m+1})\,(u_{n+1,m} - u_{n,m+1}) - (\alpha - \beta) \\ & + \varepsilon(\alpha - \beta)\left(F^{(+)}_{n+m}\, u_{n+1,m}u_{n,m+1} + F^{(-)}_{n+m}\, u_{n,m}u_{n+1,m+1}\right) = 0, \end{aligned} \tag{23a}$$

$$
\begin{aligned}
{}_rH_2^{\varepsilon}\colon\quad & (u_{n,m}-u_{n+1,m+1})(u_{n+1,m}-u_{n,m+1})+\\
&+(\beta-\alpha)(u_{n,m}+u_{n+1,m}+u_{n,m+1}+u_{n+1,m+1})-\alpha^2+\beta^2\\
&-\varepsilon\,(\beta-\alpha)^3-\varepsilon\,(\beta-\alpha)\,\left(2F^{(-)}_{n+m}u_{n,m}+2F^{(+)}_{n+m}u_{n+1,m}+\alpha+\beta\right)\cdot\\
&\cdot\left(2F^{(-)}_{n+m}u_{n+1,m+1}+2F^{(+)}_{n+m}u_{n,m+1}+\alpha+\beta\right)\,=\,0,
\end{aligned}
\tag{23b}
$$

$$
\begin{aligned}
{}_rH_3^{\varepsilon}\colon\quad & \alpha(u_{n,m}u_{n+1,m}+u_{n,m+1}u_{n+1,m+1})\\
&-\beta(u_{n,m}u_{n,m+1}+u_{n+1,m}u_{n+1,m+1})+(\alpha^2-\beta^2)\delta\\
&-\,\frac{\varepsilon(\alpha^2-\beta^2)}{\alpha\beta}\left(F^{(+)}_{n+m}\,u_{n+1,m}u_{n,m+1}+F^{(-)}_{n+m}\,u_{n,m}u_{n+1,m+1}\right)=0,
\end{aligned}
\tag{23c}
$$

Trapezoidal H^4 equations:

$$
\begin{aligned}
{}_tH_1\colon\quad & (u_{n,m}-u_{n+1,m})\,(u_{n,m+1}-u_{n+1,m+1})\,-\\
&-\alpha_2\varepsilon^2\left(F^{(+)}_{m}u_{n,m+1}u_{n+1,m+1}+F^{(-)}_{m}u_{n,m}u_{n+1,m}\right)-\alpha_2=0,
\end{aligned}
\tag{24a}
$$

$$
\begin{aligned}
{}_tH_2\colon\quad & (u_{n,m}-u_{n+1,m})\,(u_{n,m+1}-u_{n+1,m+1})\\
&+\alpha_2\,(u_{n,m}+u_{n+1,m}+u_{n,m+1}+u_{n+1,m+1})\\
&+\frac{\varepsilon\alpha_2}{2}\left(2F^{(+)}_{m}u_{n,m+1}+2\alpha_3+\alpha_2\right)\left(2F^{(+)}_{m}u_{n+1,m+1}+2\alpha_3+\alpha_2\right)\\
&+\frac{\varepsilon\alpha_2}{2}\left(2F^{(-)}_{m}u_{n,m}+2\alpha_3+\alpha_2\right)\left(2F^{(-)}_{m}u_{n+1,m}+2\alpha_3+\alpha_2\right)\\
&+(\alpha_3+\alpha_2)^2-\alpha_3^2-2\varepsilon\alpha_2\alpha_3\,(\alpha_3+\alpha_2)=0
\end{aligned}
\tag{24b}
$$

$$
\begin{aligned}
{}_tH_3\colon\quad & \alpha_2\,(u_{n,m}u_{n+1,m+1}+u_{n+1,m}u_{n,m+1})\\
&-\,(u_{n,m}u_{n,m+1}+u_{n+1,m}u_{n+1,m+1})-\alpha_3\left(\alpha_2^2-1\right)\delta^2+\\
&-\,\frac{\varepsilon^2(\alpha_2^2-1)}{\alpha_3\alpha_2}\left(F^{(+)}_{m}u_{n,m+1}u_{n+1,m+1}+F^{(-)}_{m}u_{n,m}u_{n+1,m}\right)=0,
\end{aligned}
\tag{24c}
$$

H^6 equations:

$$
\begin{aligned}
{}_1D_2\colon\quad & \left(F^{(-)}_{n+m}-\delta_1F^{(+)}_{n}F^{(-)}_{m}+\delta_2F^{(+)}_{n}F^{(+)}_{m}\right)u_{n,m}\\
&+\left(F^{(+)}_{n+m}-\delta_1F^{(-)}_{n}F^{(-)}_{m}+\delta_2F^{(-)}_{n}F^{(+)}_{m}\right)u_{n+1,m}+\\
&+\left(F^{(+)}_{n+m}-\delta_1F^{(+)}_{n}F^{(+)}_{m}+\delta_2F^{(+)}_{n}F^{(-)}_{m}\right)u_{n,m+1}\\
&+\left(F^{(-)}_{n+m}-\delta_1F^{(-)}_{n}F^{(+)}_{m}+\delta_2F^{(-)}_{n}F^{(-)}_{m}\right)u_{n+1,m+1}+\\
&+\delta_1\left(F^{(-)}_{m}u_{n,m}u_{n+1,m}+F^{(+)}_{m}u_{n,m+1}u_{n+1,m+1}\right)\\
&+F^{(+)}_{n+m}u_{n,m}u_{n+1,m+1}+F^{(-)}_{n+m}u_{n+1,m}u_{n,m+1}=0,
\end{aligned}
\tag{25a}
$$

$$
\begin{aligned}
{}_2D_2\colon\quad & \left(F_m^{(-)}-\delta_1F_n^{(+)}F_m^{(-)}+\delta_2F_n^{(+)}F_m^{(+)}-\delta_1\lambda F_n^{(-)}F_m^{(+)}\right)u_{n,m} \\
&+\left(F_m^{(-)}-\delta_1F_n^{(-)}F_m^{(-)}+\delta_2F_n^{(-)}F_m^{(+)}-\delta_1\lambda F_n^{(+)}F_m^{(+)}\right)u_{n+1,m} \\
&+\left(F_m^{(+)}-\delta_1F_n^{(+)}F_m^{(+)}+\delta_2F_n^{(+)}F_m^{(-)}-\delta_1\lambda F_n^{(-)}F_m^{(-)}\right)u_{n,m+1} \\
&+\left(F_m^{(+)}-\delta_1F_n^{(-)}F_m^{(+)}+\delta_2F_n^{(-)}F_m^{(-)}-\delta_1\lambda F_n^{(+)}F_m^{(-)}\right)u_{n+1,m+1} \\
&+\delta_1\left(F_{n+m}^{(-)}u_{n,m}u_{n+1,m+1}+F_{n+m}^{(+)}u_{n+1,m}u_{n,m+1}\right) \\
&+F_m^{(+)}u_{n,m}u_{n+1,m}+F_m^{(-)}u_{n,m+1}u_{n+1,m+1}-\delta_1\delta_2\lambda=0,
\end{aligned}
\tag{25b}
$$

$$
\begin{aligned}
{}_3D_2\colon\quad & \left(F_m^{(-)}-\delta_1F_n^{(-)}F_m^{(-)}+\delta_2F_n^{(+)}F_m^{(+)}-\delta_1\lambda F_n^{(-)}F_m^{(+)}\right)u_{n,m} \\
&+\left(F_m^{(-)}-\delta_1F_n^{(+)}F_m^{(-)}+\delta_2F_n^{(-)}F_m^{(+)}-\delta_1\lambda F_n^{(+)}F_m^{(+)}\right)u_{n+1,m} \\
&+\left(F_m^{(+)}-\delta_1F_n^{(-)}F_m^{(+)}+\delta_2F_n^{(+)}F_m^{(-)}-\delta_1\lambda F_n^{(-)}F_m^{(-)}\right)u_{n,m+1} \\
&+\left(F_m^{(+)}-\delta_1F_n^{(+)}F_m^{(+)}+\delta_2F_n^{(-)}F_m^{(-)}-\delta_1\lambda F_n^{(+)}F_m^{(-)}\right)u_{n+1,m+1} \\
&+\delta_1\left(F_n^{(-)}u_{n,m}u_{n,m+1}+F_n^{(+)}u_{n+1,m}u_{n+1,m+1}\right) \\
&+F_m^{(-)}u_{n,m+1}u_{n+1,m+1}+F_m^{(+)}u_{n,m}u_{n+1,m}-\delta_1\delta_2\lambda=0,
\end{aligned}
\tag{25c}
$$

$$
\begin{aligned}
D_3\colon\quad & F_n^{(+)}F_m^{(+)}u_{n,m}+F_n^{(-)}F_m^{(+)}u_{n+1,m}+F_n^{(+)}F_m^{(-)}u_{n,m+1} \\
&+F_n^{(-)}F_m^{(-)}u_{n+1,m+1}+F_m^{(-)}u_{n,m}u_{n+1,m} \\
&+F_n^{(-)}u_{n,m}u_{n,m+1}+F_{n+m}^{(-)}u_{n,m}u_{n+1,m+1}+ \\
&+F_{n+m}^{(+)}u_{n+1,m}u_{n,m+1}+F_n^{(+)}u_{n+1,m}u_{n+1,m+1} \\
&+F_m^{(+)}u_{n,m+1}u_{n+1,m+1}=0,
\end{aligned}
\tag{25d}
$$

$$
\begin{aligned}
{}_1D_4\colon\quad & \delta_1\left(F_n^{(-)}u_{n,m}u_{n,m+1}+F_n^{(+)}u_{n+1,m}u_{n+1,m+1}\right)+ \\
&+\delta_2\left(F_m^{(-)}u_{n,m}u_{n+1,m}+F_m^{(+)}u_{n,m+1}u_{n+1,m+1}\right)+ \\
&+u_{n,m}u_{n+1,m+1}+u_{n+1,m}u_{n,m+1}+\delta_3=0,
\end{aligned}
\tag{25e}
$$

$$
\begin{aligned}
{}_2D_4\colon\quad & \delta_1\left(F_n^{(-)}u_{n,m}u_{n,m+1}+F_n^{(+)}u_{n+1,m}u_{n+1,m+1}\right)+ \\
&+\delta_2\left(F_{n+m}^{(-)}u_{n,m}u_{n+1,m+1}+F_{n+m}^{(+)}u_{n+1,m}u_{n,m+1}\right)+ \\
&+u_{n,m}u_{n+1,m}+u_{n,m+1}u_{n+1,m+1}+\delta_3=0.
\end{aligned}
\tag{25f}
$$

B Correlation formulae between the non autonomous Q_{V} and the non autonomous YdKN in the m direction.

For the sake of completeness let us write down the non autonomous YdKN in the m direction:

$$
\frac{\mathrm{d}u_m}{\mathrm{d}t}=\frac{A_m(u_m)u_{m+1}u_{m-1}+B_m(u_m)(u_{m+1}+u_{m-1})+C_m(u_m)}{u_{m+1}-u_{m-1}}. \tag{26}
$$

Here the m-dependent coefficients are given by:

$$A_m(u_m) = au_m^2 + 2b_m u_m + c_m, \tag{27a}$$
$$B_m(u_m) = b_{m+1}u_m^2 + du_m + e_{m+1}, \tag{27b}$$
$$C_m(u_m) = c_{m+1}u_m^2 + 2e_m u_m + f, \tag{27c}$$

where b_m, c_m and e_m are 2-periodic functions, i.e.

$$b_m = b_0 + b_1(-1)^m,\ c_m = c_0 + c_1(-1)^m,\ e_m = e_0 + e_1(-1)^m. \tag{28}$$

The correlation formulae read:

$$\begin{aligned}
a &= a_1a_{5,0} - a_{2,0}^2 + \\
&\quad -a_{2,1}^2 + a_{2,2}^2 + a_{2,3}^2 + (-1)^n(2a_{2,0}a_{2,1} - 2a_{2,2}a_{2,3} + a_1a_{5,1}), \\
b_0 &= \frac{1}{2}\{a_{2,0}(a_{5,0} - a_{3,0} - a_{4,0}) + a_1a_{6,0} - a_{2,2}a_{3,2} - a_{2,3}a_{4,3} + a_{2,1}a_{5,1} + \\
&\quad +(-1)^n[a_{2,1}(a_{5,0} + a_{3,0} + a_{4,0}) + a_{2,3}a_{3,2} + a_1a_{6,1} + a_{2,0}a_{5,1} + \\
&\quad +a_{2,2}a_{4,3}]\}, \\
b_1 &= \frac{1}{2}\{a_{2,2}(a_{4,0} - a_{3,0} - a_{5,0}) - a_{2,3}a_{5,1} + a_{2,1}a_{4,3} - a_{2,0}a_{3,2} + a_1a_{6,2} + \\
&\quad +(-1)^n[a_1a_{6,3} + a_{2,3}(a_{3,0} - a_{4,0} - a_{5,0}) + a_{2,1}a_{3,2} - a_{2,0}a_{4,3} - \\
&\quad -a_{2,2}a_{5,1}]\}, \\
c_0 &= a_{2,0}a_{6,0} - a_{4,0}a_{3,0} + a_{2,1}a_{6,1} - a_{2,3}a_{6,3} - a_{2,2}a_{6,2} + \\
&\quad -(-1)^n[a_{2,2}a_{6,3} + a_{4,3}a_{3,2} + a_{2,3}a_{6,2} - a_{2,0}a_{6,1} - a_{2,1}a_{6,0}], \\
c_1 &= a_{2,1}a_{6,3} - a_{4,0}a_{3,2} + a_{2,0}a_{6,2} - a_{2,3}a_{6,1} - a_{2,2}a_{6,0} - \\
&\quad -(-1)^n[a_{2,2}a_{6,1} + a_{4,3}a_{3,0} + a_{2,3}a_{6,0} - a_{2,1}a_{6,2} - a_{2,0}a_{6,3}], \\
d &= \frac{1}{2}[a_{5,0}^2 - a_{4,0}^2 - a_{3,0}^2 + a_1a_7 + a_{3,2}^2 + a_{4,3}^2 - a_{5,1}^2 + \\
&\quad +4(-1)^n(a_{2,1}a_{6,0} + a_{2,0}a_{6,1} + a_{2,3}a_{6,2} + a_{2,2}a_{6,3})], \\
e_0 &= \frac{1}{2}\{a_{6,0}[a_{5,0} - a_{4,0} - a_{3,0}] + a_{2,0}a_7 - a_{5,1}a_{6,1} + a_{3,2}a_{6,2} + a_{4,3}a_{6,3} - \\
&\quad -(-1)^n[a_{3,2}a_{6,3} + a_{4,3}a_{6,2} + a_{5,1}a_{6,0} - a_{6,1}(a_{3,0} + a_{4,0} + a_{5,0}) - a_{2,1}a_7]\}, \\
e_1 &= \frac{1}{2}\{-a_{6,2}[a_{3,0} - a_{4,0} + a_{5,0}] + a_{3,2}a_{6,0} - a_{4,3}a_{6,1} + a_{5,1}a_{6,3} + a_{2,2}a_7 + \\
&\quad +(-1)^n[a_{4,3}a_{6,0} - a_{3,2}a_{6,1} + a_{5,1}a_{6,2} + a_{6,3}(a_{3,0} - a_{5,0} - a_{4,0}) + a_{2,3}a_7]\}, \\
f &= a_{5,0}a_7 - a_{6,0}^2 + a_{6,2}^2 + a_{6,3}^2 - a_{6,1}^2 - (-1)^n(2a_{6,2}a_{6,3} - 2a_{6,0}a_{6,1} + a_{5,1}a_7).
\end{aligned} \tag{29}$$

References

[1] V. E. Adler, A.I. Bobenko, and Yu. B. Suris. Classification of integrable equations on quad-graphs. the consistency approach. *Comm. Math. Phys.*, **233**, 513–543, 2003.

[2] V. E. Adler, A. I. Bobenko, and Yu. B. Suris. Discrete nonlinear hyperbolic equations. classification of integrable cases. *Funct. Anal. Appl.*, **43**, 3–17, 2009.

[3] M. Bellon and C. M. Viallet, Algebraic Entropy. *Comm. Math. Phys.* **204**, 1999, pp. 425437.

[4] R. Boll, Classification of 3D consistent quad-equations. *J. Nonlinear Math. Phys.*, **18**(3), 337–365, 2011.

[5] R. Boll, Corrigendum Classification of 3D consistent quad–equations, *J.Nonl. Math. Phys.* **19** (4), 1292001, 2012.

[6] R. Boll, Classification and Lagrangian structure of 3D consistent quad–equations, Ph. D. dissertation, 2012.

[7] D. K. Demskoy and C. M. Viallet, Algebraic entropy for semi-discrete equations. *J. Phys. A: Math. Theor.* **45** 2012, p. 352001.

[8] G. Gubbiotti PhD thesis to be submitted in 2016.

[9] G. Gubbiotti, C. Scimiterna and D. Levi, On quad equations consistent on the cube, submitted to *J. Phys. A: Math. Theor.* .

[10] G. Gubbiotti, C. Scimiterna and D. Levi, The non-autonomous YdKN equation and generalized symmetries of Boll equations, submitted to *J. Phys. A: Math. Theor.*.

[11] G. Gubbiotti, C. Scimiterna and D. Levi, Linearizability and fake Lax pair for a consistent around the cube nonlinear non–autonomous quad–graph equation, submitted to *Teoreticheskaya i Matematicheskaya Fizika*.

[12] D. Levi, Nonlinear differential-difference equations as Bäcklund transformations, *J. Phys. A: Math. Gen.* **14**, 1083–1098, 1981.

[13] D. Levi and R. Benguria, Bäcklund transformations and nonlinear differential difference equations, *Proc. Natl. Acad. Sci. USA* **77**, 5025–5027, 1980.

[14] D. Levi, M. Petrera, C. Scimiterna, and R. I. Yamilov. On Miura transformations and Volterra-type equations associated with the Adler–Bobenko–Suris equations. *SIGMA*, **4**, 077(14pp), 2008.

[15] D. Levi, P. Winternitz and R. I. Yamilov, Symmetries of the continuous and discrete Krichever–Novikov equation, *SIGMA*, **7**, 097 (21 pp), 2011.

[16] D. Levi and R. I. Yamilov, Classification of Evolutionary Equations on the Lattice I. The General Theory, arXiv:solv-int/9511006 v1 16 Nov 1995.

[17] D. Levi and R. I. Yamilov. Conditions for the existence of higher symmetries of the evolutionary equations on the lattice. *J. Math. Phys.*, **38**(12):6648–6674, 1997.

[18] G. Gubbiotti, and M. C. Hay, A *SymPy* module to calculate algebraic entropy for difference equations and quadrilateral partial difference equations, 2015, in preparation.

[19] J. Hietarinta, and C. M. Viallet, Searching for integrable lattice maps using factorization, *J. Phys. A: Math. Theor.* **40**, 2007, 12629–12643.

[20] O. G. Rasin and P. E. Hydon. Symmetries of integrable difference equations on the quad-graph. *Stud. Appl. Math.*, **119**, 253–269, 2007.

[21] A. Tongas, D. Tsoubelis and P. Xenitidis, Affine linear and D_4 symmetric lattice equations: symmetry analysis and reductions, *J. Phys. A: Math. Theor.* **40**, 13353–13384, 2007 arXiv:0707.3730.

[22] S. Tremblay, B. Grammaticos and A. Ramani, Integrable lattice equations and their growth properties, *Phys. Lett. A* **278** (2001) 319–324.

[23] C. M. Viallet, Algebraic *Entropy* for lattice equations, `arXiv:math-ph/0609043`, 2006.

[24] C. M. Viallet, Integrable lattice maps: Q_5 a rational version of Q_4, *Glasgow Math. J.*, **51A**, 2009 pp 157-163.

[25] C. M. Viallet, Algebraic entropy for differential-delay equations `arXiv:1408.6161`, 2014.

[26] P. D. Xenitidis, Integrability and symmetries of difference equations: the Adler-Bobenko-Suris case, Proc. 4^th Workshop Group Analysis of Differential Equations and Integrable Systems, (2009) 226–242 (arXiv:0902.3954).

[27] P. D. Xenitidis and V. G. Papageorgiou, Symmetries and integrability of discrete equations defined on a black and white lattice, *J. Phys. A: Math. Theor.*, **42** 454025 (13 pp.), 2009.

[28] R. I. Yamilov Classification of discrete evolution equations *Uspekhi Mat. Nauk* **38** 155–156, 1983 (in Russian).

[29] R. I. Yamilov, *Discrete equations of the form* $\dot{u}_n = F(u_{n-1}, u_n, u_{n+1})$ $n \in Z$ *with an infinite number of local conservation laws* Dissertation for Candidate of Science (Ufa: Soviet Academy of Sciences) 1984 (in Russian).

[30] R. I. Yamilov, Symmetries as integrability criteria for differential difference equations, *J.* Phys. A: Math. Gen. **39** R541–R623, 2006.